



## Evidence of human-to-dog transmission of monkeypox virus

Published Online  
August 10, 2022  
[https://doi.org/10.1016/S0140-6736\(22\)01487-8](https://doi.org/10.1016/S0140-6736(22)01487-8)

See Online for appendix

Human monkeypox virus is spreading in Europe and the USA among individuals who have not travelled to endemic areas.<sup>1</sup> On July 23, 2022, monkeypox was declared a Public Health Emergency of International Concern by WHO Director-General Tedros Adhanom Ghebreyesus.<sup>2</sup> Human-to-human transmission of monkeypox virus usually occurs through close contact with the lesions, body fluids, and respiratory droplets of infected people or animals.<sup>3</sup> The possibility of sexual transmission is being investigated, as the current outbreak appears to be concentrated in men who have sex with men and has been associated with unexpected anal and genital lesions.<sup>1,4</sup> Whether

domesticated cats and dogs could be a vector for monkeypox virus is unknown. Here we describe the first case of a dog with confirmed monkeypox virus infection that might have been acquired through human transmission.


Two men who have sex with men attended Pitié-Salpêtrière Hospital, Paris, France, on June 10, 2022 (appendix). One man (referred to as patient 1 going forward) is Latino, aged 44 years, and lives with HIV with undetectable viral loads on antiretrovirals; the second man (patient 2) is White, aged 27 years, and HIV-negative. The men are non-exclusive partners living in the same household. They each signed a consent form for the use of their clinical and biological data, and for the publication of anonymised photographs. The men had presented with anal ulceration 6 days after sex with other partners. In

patient 1, anal ulceration was followed by a vesiculopustular rash on the face, ears, and legs; in patient 2, on the legs and back (figure A, B). In both cases, rash was associated with asthenia, headaches, and fever 4 days later.

Monkeypox virus was assayed by real-time PCR (LightCycler 480 System; Roche Diagnostics, Meylan, France). In patient 1, virus was detected in skin and oropharynx samples; whereas in patient 2, virus was detected in anal and oropharynx samples.

12 days after symptom onset, their male Italian greyhound, aged 4 years and with no previous medical disorders, presented with mucocutaneous lesions, including abdomen pustules and a thin anal ulceration (figure C, D; appendix). The dog tested positive for monkeypox virus by use of a PCR protocol adapted from Li and colleagues<sup>5</sup> that involved scraping skin lesions and swabbing the anus and oral cavity. Monkeypox virus DNA sequences from the dog and patient 1 were compared by next-generation sequencing (MinION; Oxford Nanopore Technologies, Oxford, UK). Both samples contained virus of the hMPXV-1 clade, lineage B.1, which has been spreading in non-endemic countries since April, 2022, and, as of Aug 4, 2022, has infected more than 1700 people in France, mostly concentrated in Paris, where the dog first developed symptoms. Moreover, the virus that infected patient 1 and the virus that infected the dog showed 100% sequence homology on the 19.5 kilobase pairs sequenced. The men reported co-sleeping with their dog. They had been careful to prevent their dog from contact with other pets or humans from the onset of their own symptoms (ie, 13 days before the dog started to present cutaneous manifestations).

In endemic countries, only wild animals (rodents and primates) have been found to carry monkeypox virus.<sup>6</sup> However, transmission of monkeypox virus in prairie dogs has been described



**Figure:** Skin and mucosal lesions in two male patients and their dog with confirmed monkeypox virus (A) Pustular lesion of the thigh, with central umbilication and the onset of necrosis, in patient 1. (B) Erosive and pustular anal lesions in patient 2. (C) Two slightly crusty erythematous papules in the dog. (D) Millimetric erosive anal lesion in the dog.

For monkeypox cases in France  
see <https://www.santepubliquefrance.fr/les-actualites/2022/cas-de-variole-du-singe-point-de-situation-au-4-aout-2022>

in the USA<sup>7</sup> and in captive primates in Europe<sup>8</sup> that were in contact with imported infected animals. Infection among domesticated animals, such as dogs and cats, has never been reported.

To the best of our knowledge, the kinetics of symptom onset in both patients and, subsequently, in their dog suggest human-to-dog transmission of monkeypox virus. Given the dog's skin and mucosal lesions as well as the positive monkeypox virus PCR results from anal and oral swabs, we hypothesise a real canine disease, not a simple carriage of the virus by close contact with humans or airborne transmission (or both). Our findings should prompt debate on the need to isolate pets from monkeypox virus-positive individuals. We call for further investigation on secondary transmissions via pets.

We declare no competing interests. This work was supported by the French National Research Agency on HIV/AIDS, Hepatitis and Emerging Infectious Diseases. We thank the patients and all the clinical and technical staff of the Infectious Diseases and Virology Departments of Pitié-Salpêtrière Hospital (especially Anne-Geneviève Marcelin and Vincent Calvez) and Bichat-Claude Bernard Hospital, who provided care for the patients and conducted the virological explorations.

\*Sophie Seang, Sonia Burrel, Eve Todesco, Valentin Leducq, Gentine Monsel, Diane Le Pluart, Christophe Cordevant, Valérie Pourcher, Romain Palich  
sophie.seang@aphp.fr

Infectious Diseases Department (SS, GM, VP, RP) and Virology Department (SB, ET, VL), Sorbonne University, Pitié-Salpêtrière Hospital, AP-HP, Pierre Louis Epidemiology and Public Health institute, INSERM 1136, Paris 75013, France; Infectious Diseases Department, Bichat-Claude Bernard University Hospital, AP-HP, Paris, France (DLP); Strategy and Programmes Department, Research and Reference Division, Anses, Maisons-Alfort, France (CC)

- 1 Thornhill JP, Barkati S, Walmsley S, et al. Monkeypox virus infection in humans across 16 countries—April–June 2022. *N Engl J Med* 2022; published online July 21. <https://doi.org/10.1056/NEJMoa2207323>.
- 2 Wenham C, Eccleston-Turner M. Monkeypox as a PHEIC: implications for global health governance. *Lancet* 2022; published online Aug 1. [https://doi.org/10.1016/S0140-6736\(22\)01437-4](https://doi.org/10.1016/S0140-6736(22)01437-4).
- 3 Brown K, Leggat PA. Human monkeypox: current state of knowledge and implications for the future. *Trop Med Infect Dis* 2016; **1**: e8.

- 4 Tarín-Vicente EJ, Alemany A, Agud-Dios M, et al. Clinical presentation and virological assessment of confirmed human monkeypox virus cases in Spain: a prospective observational cohort study. *Lancet* 2022; published online Aug 8. [https://doi.org/10.1016/S0140-6736\(22\)01436-2](https://doi.org/10.1016/S0140-6736(22)01436-2).
- 5 Li Y, Zhao H, Wilkins K, Hughes C, Damon IK. Real-time PCR assays for the specific detection of monkeypox virus West African and Congo Basin strain DNA. *J Virol Methods* 2010; **169**: 223–27.
- 6 Khodakevich L, Jezek Z, Kinzanzka K. Isolation of monkeypox virus from wild squirrel infected in nature. *Lancet* 1986; **1**: 98–99.
- 7 Parker S, Buller RM. A review of experimental and natural infections of animals with monkeypox virus between 1958 and 2012. *Future Virol* 2013; **8**: 129–57.
- 8 Arita I, Henderson DA. Smallpox and monkeypox in non-human primates. *Bull World Health Organ* 1968; **39**: 277–83.

## WHO's Palestinian statistics: what's Israel got to do with it?

We are writing to express our serious concerns and surprise over some of the statements and quotations offered in Richard Horton's Offline<sup>1</sup> about WHO's decision not to publish health statistics for the West Bank and the Gaza Strip in their *World Health Statistics 2022*.<sup>2</sup> In fact, we entirely agree with Horton's central point about the need to publish accurate health statistics from all jurisdictions. We, too, cannot understand why Palestinian health data were not included, especially as Palestine is a member of the WHO Eastern Mediterranean Regional Office group. Moreover, we are deeply concerned by the health conditions and status of Palestinians, as reported by the Director-General in May, 2022.<sup>3</sup>

That being said, we are perplexed by Horton's conflation of WHO's omission with various purported sins committed over more than 100 years, first by the Zionists, then perpetuated by Israel. We are confounded by this non sequitur since Israel could not, and did not, have any role in WHO's decision. Why bring Israel into the discussion at all? Both the content and tone strike us simply as a gratuitous

swipe at Israel, Zionism, and Jews, which can have no constructive role in advancing Palestinian health. A few examples we hope will elucidate our concerns.

Genocide is a highly charged word that can push many buttons. In this context we are all too familiar with the preposterous and poisonous claim that the Jews or just the Zionists are somehow out to destroy the Palestinian people. The complex history of our region relates numerous Israeli acceptances of international decisions and attempts at reconciliation with Palestinian authorities and leadership, which, tragically, were met with responses aimed at eliminating the Jewish people in Israel.<sup>4</sup> Use of the loaded term genocide also, sadly, feeds directly into a narrative of some who accuse Israel and its supporters of responsibility for a Palestinian Holocaust—a mirror image of what was perpetrated by Nazi Germany and its European collaborators on the Jews.

Taiwan, with 20 million inhabitants, is also, unhelpfully, not included in WHO's health statistics.<sup>2</sup> Taiwan's effort to join WHO as an observer failed once again earlier this year.<sup>5</sup> Clearly, Taiwan is a fully-fledged sovereign country, whatever the Chinese leadership thinks or claims. Although worthy of criticism, is WHO's exclusion of Taiwan's membership a form of genocide? To us, and surely to other fair-minded readers of *The Lancet*, the abuse of this word in this context points to another, not so subtle, attempt to serve an agenda of delegitimising Israel.

To claim that "life expectancy, infant and under-5 mortality, and maternal mortality were all worse than those of the occupying nation",<sup>1</sup> which Horton attributes to Rik Peeperkorn of WHO, does not take into account the complexity of the multifactorial reasons for the tragic state of Palestinian health. Without wishing to diminish Israel's contribution to the difficult state of Palestinian health at times, it is clear that a major part of the



Published Online  
August 18, 2022  
[https://doi.org/10.1016/S0140-6736\(22\)01583-5](https://doi.org/10.1016/S0140-6736(22)01583-5)